

Clinical Policy: Betamethasone Dipropionate Spray (Sernivo)

Reference Number: CP.PMN.182

Effective Date: 12.01.18 Last Review Date: 11.19

Line of Business: Commercial, Medicaid Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Betamethasone dipropionate 0.05% spray (Sernivo®) is a topical corticosteroid.

FDA Approved Indication(s)

Sernivo is indicated for the treatment of mild to moderate plaque psoriasis (PsO) in patients 18 years of age or older.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Sernivo is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Plaque Psoriasis (must meet all):

- 1. Diagnosis of PsO;
- 2. Age \geq 18 years;
- 3. Failure of a medium to ultra high potency topical corticosteroid (*see Appendix B*), unless contraindicated or clinically significant adverse effects are experienced;
- 4. Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: calcipotriene, calcitriol, or tazarotene;
- 5. Dose does not exceed 120 mL every 4 weeks.

Approval duration: 1 month

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Plaque Psoriasis (must meet all):

- 1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
- 2. Member is responding positively to therapy;
- 3. If request is for a dose increase, new dose does not exceed 120 mL every 4 weeks.



Approval duration: Up to 1 month of total treatment (a single continuous course of therapy up to 4 weeks is recommended)

B. Other diagnoses/indications (must meet 1 or 2):

- 1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
 - Approval duration: Duration of request or 12 months (whichever is less); or
- 2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial and CP.PMN.53 for Medicaid.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key FDA: Food and Drug Administration

PsO: plaque psoriasis

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

| Drug Name | Dosing Regimen | Dose Limit/ Maximum Dose |
|--|---------------------------------|-----------------------------|
| calcipotriene (Dovonex®) | Apply topically to the affected | 100 g/week |
| cream, ointment, solution | area(s) BID | |
| calcitriol (Vectical TM) | Apply topically to the affected | 200 g/week |
| ointment | area(s) BID | |
| tazarotene (Tazorac®) gel, | Apply topically to the | Once daily application |
| cream | affected area(s) QHS | |
| Ultra High Potency Topical C | Corticosteroids | |
| augmented betamethasone | Apply topically to the affected | Should not be used for |
| dipropionate 0.05% | area(s) BID | longer than 2 |
| (Diprolene [®] , Alphatrex [®]) | | consecutive weeks |
| ointment, gel | | |
| clobetasol propionate 0.05% | | |
| (Temovate [®] , Temovate E [®]) | | |
| cream, ointment, gel, solution | | |
| diflorasone diacetate 0.05% | | |
| (Apexicon®) ointment | | |
| halobetasol propionate 0.05% | | |
| (Ultravate®) cream, ointment | | |



| Drug Name | Dosing Regimen | Dose Limit/ | | | | |
|--|---------------------------------|------------------------|--|--|--|--|
| | | Maximum Dose | | | | |
| High Potency Topical Corticosteroids | | | | | | |
| augmented betamethasone | Apply topically to the affected | Should not be used for | | | | |
| dipropionate 0.05% | area(s) BID | longer than 2 | | | | |
| (Diprolone [®] , Diprolene [®] AF) | | consecutive weeks | | | | |
| cream, lotion | | | | | | |
| betamethasone dipropionate | | | | | | |
| 0.05% ointment | | | | | | |
| desoximetasone (Topicort®) | | | | | | |
| 0.25%, 0.05% cream, | | | | | | |
| ointment, gel | | | | | | |
| diflorasone 0.05% (Apexicon | | | | | | |
| E [®]) cream | | | | | | |
| fluocinonide acetonide 0.05% | | | | | | |
| cream, ointment, gel, solution | | | | | | |
| triamcinolone acetonide 0.5% | | | | | | |
| (Aristocort [®] , Kenalog [®]) | | | | | | |
| cream, ointment | | | | | | |
| Medium/Medium to High Pot | | | | | | |
| betamethasone dipropionate | Apply topically to the affected | Should not be used for | | | | |
| 0.05% cream | area(s) BID | longer than 2 | | | | |
| desoximetasone 0.05% | | consecutive weeks | | | | |
| (Topicort®) cream, ointment, | | | | | | |
| gel | | | | | | |
| fluocinolone acetonide | | | | | | |
| 0.025% (Synalar®) cream, | | | | | | |
| ointment | | | | | | |
| fluticasone propionate 0.05% | | | | | | |
| (Cutivate [®]) cream | | | | | | |
| mometasone furoate 0.1% | | | | | | |
| (Elocon®) cream, lotion, | | | | | | |
| ointment | | | | | | |
| triamcinolone acetonide | | | | | | |
| 0.1%, 0.25%, 0.5% | | | | | | |
| (Aristocort®, Kenalog®) | | | | | | |
| cream, ointment | | | | | | |

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings None reported



V. Dosage and Administration

| Drug Name | Dosing Regimen | Maximum Dose |
|----------------------------|--------------------------------|----------------|
| Betamethasone dipropionate | Apply spray topically to | Not applicable |
| 0.05% (Sernivo) | affected areas BID for up to 4 | |
| | weeks. Avoid use on face, | |
| | scalp, axilla, groin, or other | |
| | intertriginous areas. | |

VI. Product Availability

Spray: 60 mL, 120 mL

VII. References

- 1. Sernivo Prescribing Information. San Antonio, TX: DPT Laboratories; November 2018. Available at: http://www.sernivo.com/. Accessed August 6, 2019.
- 2. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 3. Guidelines of care for the management and treatment of psoriasis with topical therapies. J Am Acad Dermatol 2009 Apr;60(4):643-59.
- 3. DRUGDEX® System [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed August 6, 2019.

| Reviews, Revisions, and Approvals | Date | P&T Approval Date |
|---|----------|-------------------------|
| Policy created: adapted from previously approved policy CP.CPA.255 (retired); age requirement added; no significant changes; references reviewed and updated. | 08.14.18 | 11.18 |
| 4Q 2019 annual review: no significant changes; references reviewed and updated. | 08.08.19 | 11.19 |

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and



limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.